

Suicide worldwide in 2021

Global health estimates



World Health
Organization

Suicide worldwide in 2021

Global health estimates



**World Health
Organization**

Suicide worldwide in 2021: global health estimates

ISBN 978-92-4-011006-9 (electronic version)

ISBN 978-92-4-011007-6 (print version)

© World Health Organization 2025

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Under the terms of this licence, you may copy, redistribute and adapt the work for non-commercial purposes, provided the work is appropriately cited, as indicated below. In any use of this work, there should be no suggestion that WHO endorses any specific organization, products or services. The use of the WHO logo is not permitted. If you adapt the work, then you must license your work under the same or equivalent Creative Commons licence. If you create a translation of this work, you should add the following disclaimer along with the suggested citation: “This translation was not created by the World Health Organization (WHO). WHO is not responsible for the content or accuracy of this translation. The original English edition shall be the binding and authentic edition”.

Any mediation relating to disputes arising under the licence shall be conducted in accordance with the mediation rules of the World Intellectual Property Organization (<http://www.wipo.int/amc/en/mediation/rules/>).

Suggested citation. Suicide worldwide in 2021: global health estimates. Geneva: World Health Organization; 2025. Licence: [CC BY-NC-SA 3.0 IGO](https://creativecommons.org/licenses/by-nc-sa/3.0/igo).

Cataloguing-in-Publication (CIP) data. CIP data are available at <https://iris.who.int/>.

Sales, rights and licensing. To purchase WHO publications, see <https://www.who.int/publications/book-orders>. To submit requests for commercial use and queries on rights and licensing, see <https://www.who.int/copyright>.

Third-party materials. If you wish to reuse material from this work that is attributed to a third party, such as tables, figures or images, it is your responsibility to determine whether permission is needed for that reuse and to obtain permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

General disclaimers. The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of WHO concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by WHO in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by WHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall WHO be liable for damages arising from its use.

Design and layout by Studio FFFOG.

Contents

Acknowledgements	iv
Introduction	1
Data sources and methods	2
Reasons for differences from official national estimates	2
Reasons for changes in suicide estimates compared with previous WHO publications	3
Global epidemiology of suicide	4
Regional epidemiology of suicide	9
Changes in suicide rates over time	11
Data quality	13
Conclusions	15
References	16
Annex 1	18
Annex 2	38

Acknowledgements

Guidance and oversight

Dévora Kestel and Mark van Ommeren (WHO Department of Mental Health, Brain Health and Substance Use)

Project coordination, writing and editing

Alexandra Fleischmann, Piamee Bandara, Sandersan Onie (WHO Department of Mental Health, Brain Health and Substance Use) and Elise Paul (University College London, United Kingdom of Great Britain and Northern Ireland)

Global Health Estimates and maps

Bochen Cao, Jessica Ho and Wahyu Retno Mahanani (WHO Department of Data and Analytics); Doris Ma Fat, Zoe Brillantes and Kathleen Friar (WHO Department of Surveys, CRVS, and Health Service Data)

Introduction

Suicide represents a critical public health challenge. Globally, 727 000 people died by suicide in 2021. Suicide is among the leading causes of death worldwide, with more deaths from suicide than from malaria, HIV/AIDS, breast cancer, or war and homicide. Suicide accounted for more than one in every 100 deaths (1.1%) in 2021.

Reducing suicide is an indicator in the United Nations Sustainable Development Goals (UN SDGs) under target 3.4, and it is the only mental health indicator (3.4.2). It is also a World Health Organization (WHO) priority, included as a global target in WHO's overall work plans from 2019 to 2028 (1, 2), and its *Comprehensive Mental Health Action Plan 2013–2030* (3). A comprehensive and coordinated response to suicide prevention is critical to save lives and reduce the impact on families and communities.

Keeping up-to-date records and constantly monitoring suicide is essential for effective national suicide prevention strategies and measures (4, 5). Countries should break down suicide data, at least by sex, age and method. Such disaggregated data helps to: identify which groups are most at risk, understand the scope of the problem, tailor interventions to meet the needs of specific populations, and adapt approaches to evolving trends.

Data sources and methods

This publication offers the latest data on suicide from WHO's Global Health Estimates for 2000–2021 (6). To facilitate comparisons across countries, rates were age-standardized according to the WHO World Standard Population, which assumes one standard age distribution of the population in all countries.

Data are presented globally, regionally, and by income level, broken down by age and sex over time. Country- and age-specific estimates are in Annex 1 and Annex 2.

Details of methods, data sources and analyses can be found on the WHO Global Health Estimates website (7).¹ Global health estimates for suicide were calculated using country mortality data reported to the WHO Mortality Database (8). For countries lacking comprehensive death registration or other suicide data, data from the Institute of Health Metrics and Evaluation's Global Burden of Disease 2021 were used (9). Estimates were made for 183 WHO Member States with populations greater than 90 000 in 2021. The 11 WHO Member States excluded are: Andorra, Cook Islands, Dominica, Marshall Islands, Monaco, Nauru, Niue, Palau, Saint Kitts and Nevis, San Marino, and Tuvalu.

Reasons for differences from official national estimates

The Global Health Estimates represent WHO's best estimates, based on evidence available until May 2024, rather than the official estimates of Member States, and have not necessarily been endorsed by Member States. They have been computed using standard categories, definitions and methods to ensure cross-national comparability and may not be the same as official national estimates produced using alternative, potentially equally rigorous, methods. The process involved extracting codes X60–X84 and Y870 for suicide from the WHO Mortality Database, redistributing deaths of unknown sex/age and deaths assigned to ill-defined codes, interpolating/extrapolating number of deaths for missing years, scaling total deaths by age and sex to WHO all-cause envelopes for 2000–2021, and using population estimates from the UN Population Division.

¹ As the availability and quality of causes of death data used as inputs in WHO's Global Health Estimates analysis for the years 2020 and 2021 were affected in many countries by disruptions associated with the COVID-19 pandemic, the estimates for suicide deaths in 2020 and 2021 are probably underestimated and the trends discussed here need to be interpreted with caution.


Reasons for changes in suicide estimates compared with previous WHO publications

WHO regularly updates its entire set of global health estimates, including suicide estimates, from the year 2000 on. The latest estimates for 2000–2021 used new data, drew on results of the 2021 Global Burden of Diseases study, and had substantial revisions to the methods used. This means they are not comparable with previously published estimates for 2000–2019 (10) or with earlier versions published by WHO. The suicide estimates in this publication replace all those previously published by WHO, and any differences with previously published versions should not be interpreted as time trends (11). To identify time trends, please see the latest published version of WHO Global Health Estimates (Fig. 10).



Global epidemiology of suicide

In 2021, an estimated 727 000 people died by suicide. The global age-standardized suicide rate was 8.9 per 100 000 population for 2021. Rates varied between countries from fewer than one death by suicide per 100 000 to almost 40 per 100 000 (Fig. 1).

Fig. 1. Age-standardized suicide rates (per 100 000 population), both sexes, 2021




The global age-standardized suicide rate was higher in males (12.3 per 100 000) than in females (5.6 per 100 000) (Fig. 2 and Fig. 3). For females, the highest suicide rates in individual countries reached to 20.3 per 100 000 (Fig. 2); for males they rose above 50.0 per 100 000 (Fig. 3).

Fig. 2. Age-standardized suicide rates (per 100 000 population), females, 2021**Fig. 3. Age-standardized suicide rates (per 100 000 population), males, 2021**


Globally, the age-standardized suicide rate was 2.2 times higher in males than in females. Male to female suicide ratios greater than one indicate that suicide rates are higher in males than in females. This ratio was 3.2 in high-income countries, 3.0 in low-income countries, 2.3 in upper-middle-income countries, and 1.7 in lower-middle-income countries (Fig. 4).

Fig. 4. Male to female ratio of age-standardized suicide rates, 2021



Globally, most suicides were in low- and middle-income countries (73%), where most of the world's population lives (Fig. 5). More than half of global suicides (56%) happened before the age of 50 years. Most adolescents (10-19 years) who died by suicide (84%) were from low- and middle-income countries.


Fig. 5. Global suicides, by age and country income level* (thousands), 2021



*World Bank income groups, 2024


Suicide was the third leading cause of death in young people aged 15–29 years for both sexes, after road injury and interpersonal violence (Fig. 6). For females, suicide was the second, and for males, the third leading cause of death in this age group.

Fig. 6. Top four causes of death, ages 15–29 years, 2021



Suicide was the fourth leading cause of death in 15-19-year-olds for both sexes, with the number of deaths relatively similar between females and males in this age group (Fig. 7). Suicide was the second leading cause of death in 15-19-year-old girls (after maternal conditions) and the fourth leading cause of death in males in this age group.

Fig. 7. Top four causes of death, ages 15-19 years, 2021




Regional epidemiology of suicide

Differences in age-standardized suicide rates were seen across WHO regions in 2021 (Fig. 8). Suicide rates were higher than the global average of 8.9 per 100 000 in WHO's African Region (11.5 per 100 000), European Region (10.1 per 100 000), South-East Asia Region (10.1 per 100 000) and Region of the Americas (9.2 per 100 000). The lowest suicide rate was in the Eastern Mediterranean Region (4.0 per 100 000). The rate in the Western Pacific Region was 7.5 per 100 000 in 2021.


The South-East Asia Region had a much higher female age-standardized suicide rate (8.3 per 100 000) than the global female average (5.6 per 100 000) in 2021. For males, WHO's African Region (18.4 per 100 000), Region of the Americas (14.6 per 100 000) and European Region (16.3 per 100 000) all had higher suicide rates than the global average (12.3 per 100 000).

Fig. 8. Age-standardized suicide rates (per 100 000 population) by WHO region, 2021



In 2021, most suicides happened in low- and middle-income countries (73%). Suicide rates varied slightly by income group. High-income countries had the highest age-standardized suicide rate (11.8 per 100 000). Low-income countries had a slightly lower rate (10.5 per 100 000), followed by lower-middle-income and upper-middle-income countries (9.4 per 100 000 and 7.0 per 100 000 respectively) (Fig. 9). The highest suicide rate for females was in lower-middle-income countries (7.1 per 100 000); for males, it was in high-income countries (18.1 per 100 000).

Fig. 9. Age-standardized suicide rates (per 100 000 population) by income level*, 2021




*World Bank income groups, 2024

Changes in suicide rates over time

Between 2000 and 2021, the global age-standardized suicide rate dropped by 35%. The decreases varied by WHO region: 3% in Africa, 26% in South-East Asia, 30% in the Eastern Mediterranean, 48% in Europe and 50% in the Western Pacific (Fig. 10). The only region with an increase was the Americas, where the age-standardized suicide rate rose by 17% in the same time period. The global rate also decreased for age-group specific suicide rates (i.e. 15-29 years, 30-49 years, 50-69 years, and 70+ years) in the same time period.

Fig. 10. Age-standardized suicide rates (per 100 000 population) over time by WHO region, both sexes



Progress in reducing the suicide mortality rate needs to be accelerated if the world is to meet its UN SDG target of a one-third reduction by 2030. Based on trends from 2015 to 2019, and assuming that the rate of change remains the same, only a 12% reduction will be achieved by 2030. To reach the one-third reduction, countries need to triple their efforts, increasing the annual average reduction from 1% to 3%.

A multisectoral, public health approach to suicide prevention will be critical to success. WHO's LIVE LIFE initiative for suicide prevention (12) provides a starting point for governments and communities to contribute to suicide prevention by implementing four key evidence-based interventions.

Limit access to the means of suicide

This intervention is proven to reduce suicides at a population level. It includes banning highly hazardous pesticides, restricting access to firearms, and installing barriers in public places. These measures can save many lives (13). It is estimated that in the Americas alone, restricting access to pesticides and firearms could prevent more than 120 000 suicide deaths in a decade (14). Worldwide, 15–20% of suicides involve pesticide self-poisoning. Phasing out highly hazardous pesticides (15) in countries where they are often used for suicide could greatly contribute to the goal of reducing suicides by one-third. Banning these pesticides is also a very cost-effective way of preventing suicides (16).

Interact with the media for responsible reporting of suicide

Media reports about suicide can increase suicides through imitation, especially if the report involves celebrities or describes the method used (17). In contrast, stories of survival, hope and recovery can help reduce suicides (18). WHO's Preventing suicide: a resource for media professionals (19) provides guidance on how to report on suicide responsibly. Working with national media bodies, local media outlets, and social media platforms is critical to ensure national guidelines and standards for safe reporting are adopted.

Foster socio-emotional life skills in young people

This intervention is crucial because suicide is a leading cause of death among young people globally. Adolescence is a critical time for acquiring social and emotional life skills, but it is also when mental health conditions often begin. Teaching young people socio-emotional skills, for example through school-based programmes, has been shown to reduce suicidal behaviour (20) and is a cost-effective way to prevent suicide. WHO's Helping Adolescents Thrive (HAT) (21) guidelines focus on promoting mental well-being and

preventing mental health conditions and self-harm in adolescents. Schools are encouraged to provide a safe school environment, connect students to support services, and have clear protocols for staff when suicide risk is identified. It is also important for parents to have the skills and knowledge to address mental health and risk factors.

Early identify, assess, manage and follow up anyone affected by suicidal behaviours

People at risk of suicide, including those who have self-harmed or lost someone to suicide, need proper support and care. Follow-up care and psychosocial support after self-harm have been shown to reduce the chances of it happening again (22). Health care workers and community members who might encounter at-risk individuals are well-positioned to identify and follow up with them. Health care facilities and systems should also prioritize suicide prevention to enable early intervention when people seek help or are identified as at risk. WHO's mhGAP intervention guide (mhGAP-IG) (23) provides guidelines for identifying, assessing, managing and following up on mental health conditions, including a specific module on suicide and self-harm.

To support the implementation of these key interventions, LIVE LIFE is built on six foundational cross-cutting pillars: situation analysis, multisectoral collaboration, awareness-raising and advocacy, capacity building, financing, and surveillance.

The initiative emphasizes that suicide prevention is not just the health sector's responsibility. Rather, a whole-of-government and whole-of-society approach is needed. Such an approach should include addressing the broader social conditions that shape people's daily lives and health.

Data quality

Only 60 of the 183 WHO Member States for which estimates were made for 2000-2021 had high-quality vital registration data (Fig. 11). Modelling methods were used to generate estimates for most of the others, which were mostly low- and middle-income countries.

The type and complexity of models for global health estimates vary widely across research/institutional groups and health estimates. More complex models yield more accurate uncertainty intervals. When high-quality data are available, estimates from different institutions usually agree. Discrepancies are more likely to arise for countries with low-quality, sparse, or potentially biased data. These discrepancies are best addressed by improving the primary data.


Suicide is a stigmatized cause of death and so it is often under-reported. The accuracy of death certificates and variability in coding practices need to be addressed. Strengthening country health information systems, including vital registration, is crucial to provide a more solid empirical basis for monitoring global, regional, national and subnational trends and responding to population needs.

To improve the monitoring of suicide, self-harm and associated factors, health information systems should focus on:

- strengthening death registration through civil registration and vital statistics systems, local health and demographic studies and other sources;
- enhancing cause-of-death data collection through vital registration and verbal autopsies in communities;
- conducting regular household health surveys; and
- ensuring complete facility recording and reporting with regular quality control.

Most suicides occur in low- and middle-income countries, where good-quality vital registration data are urgently needed. Improving suicide surveillance is important to inform planning, priority setting, monitoring and evaluation in countries, and to accurately assess progress towards global suicide mortality targets.

Fig. 11. Data quality of suicide estimates, 2021



Conclusions

The suicide mortality data in this publication underscore the urgent need for action to prevent suicide. Suicide is a global public health issue affecting all ages, sexes and regions of the world. Each data point here represents a life lost to suicide; and each loss is one too many.

While the global age-standardized suicide rate is declining overall, this trend is not seen in all countries. If the decline continues at its current rate, global targets to reduce suicide by one third by 2030 will not be met.

Lives will be lost, even though suicides are preventable.

The LIVE LIFE implementation guide for suicide prevention outlines four effective evidence-based interventions to prevent suicide. These are:

- limiting access to the means of suicide, such as highly hazardous pesticides and firearms;
- interacting with the media for responsible reporting of suicide;
- fostering socio-emotional life skills in adolescents; and
- early identification, assessment, management and follow-up of anyone who is affected by suicidal behaviours.

Further action, strengthening and acceleration of ongoing suicide prevention efforts are needed to save lives.

References

1. Seventy-first World Health Assembly: Agenda item 11.1, 25 May 2018: Thirteenth General Programme of Work, 2019–2023. Geneva: World Health Organization; 2018 (WHA71.1; <https://iris.who.int/handle/10665/279451>, accessed 20 February 2025).
2. A global health strategy for 2025–2028: advancing equity and resilience in a turbulent world: fourteenth General Programme of Work. Geneva: World Health Organization; 2025 (<https://iris.who.int/handle/10665/380456>, accessed 20 February 2025). Licence: CC BY-NC-SA 3.0 IGO.
3. Comprehensive mental health action plan 2013–2030. Geneva: World Health Organization; 2021 (<https://iris.who.int/handle/10665/345301>, accessed 20 February 2025). Licence: CC BY-NC-SA 3.0 IGO.
4. Preventing suicide: a global imperative. Geneva: World Health Organization; 2014. (<https://iris.who.int/handle/10665/131056>, accessed 20 February 2025).
5. LIVE LIFE: an implementation guide for suicide prevention in countries. Geneva: World Health Organization; 2021 (<https://iris.who.int/handle/10665/341726>, accessed 20 February 2025). Licence: CC BY-NC-SA 3.0 IGO.
6. Global Health Estimates [online database]. Geneva: World Health Organization; 2025 (<https://www.who.int/data/global-health-estimates>, accessed 20 February 2025).
7. WHO methods and data sources for country-level causes of death 2000–2021. Geneva: World Health Organization; 2024 (https://cdn.who.int/media/docs/default-source/gho-documents/global-health-estimates/ghe2021_cod_methods.pdf?sfvrsn=dca346b7_1, accessed 20 February 2025).
8. WHO mortality database [online database]. Geneva: World Health Organization; 2025 (<https://www.who.int/data/data-collection-tools/who-mortality-database>, accessed 20 February 2025).
9. 2021 Global Burden of Diseases (GBD) [online database]. Seattle: Institute for Health Metrics and Evaluation; 2024 (<https://vizhub.healthdata.org/gbd-results>, accessed 20 February 2025).
10. Suicide worldwide in 2019: Global Health Estimates. Geneva: World Health Organization; 2021. <https://iris.who.int/handle/10665/341728>, accessed 20 February 2025). Licence: CC BY-NC-SA 3.0 IGO.
11. Reasons for changes in GHE estimates in this revision. In: WHO methods and data sources for country-level causes of death 2000–2021. Geneva: World Health Organization; 2024 (https://cdn.who.int/media/docs/default-source/gho-documents/global-health-estimates/ghe2021_cod_methods.pdf?sfvrsn=dca346b7_1, accessed 20 February 2025).
12. WHO LIVE LIFE initiative [website]. Geneva: World Health Organization (<https://www.who.int/initiatives/live-life-initiative-for-suicide-prevention>, accessed 20 February 2025).
13. LIVE LIFE: an implementation guide for suicide prevention in countries. Geneva: World Health Organization; 2021 (<https://iris.who.int/handle/10665/341726>, accessed 20 February 2025). Licence: CC BY-NC-SA 3.0 IGO.
14. Lange S, Kim KV, Jiang H, Shield KD, Rehm J, Hennis AJM et al. Forecasting the impact of means restriction on the suicide mortality rate in the Region of the Americas: an ecological modeling study. *Lancet Reg Health Am*. 2024;36:100831 (<https://doi.org/10.1016/j.lana.2024.100831>, accessed 20 February 2025).

15. Preventing suicide by phasing out highly hazardous pesticides. Geneva: World Health Organization; 2024 (<https://www.who.int/publications/m/item/preventing-suicide-by-phasing-out-highly-hazardous-pesticides>, accessed 20 February 2025).
16. WHO menu of cost-effective interventions for mental health. Geneva: World Health Organization; 2021 (<https://iris.who.int/handle/10665/343074>, accessed 20 February 2025). Licence: CC BY-NC-SA 3.0 IGO.
17. Niederkrotenthaler T, Braun M, Pirkis J, Till B, Stack S, Sinyor M et al. Association between suicide reporting in the media and suicide: systematic review and meta-analysis. *BMJ*. 2020;368:m575 (<https://doi.org/10.1136/bmj.m575>, accessed 20 February 2025).
18. Niederkrotenthaler T, Tran U, Gould M, Sinyor M, Sumner S, Strauss MJ et al. Association of Logic's hip hop song 1-800-273-8255 with lifeline calls and suicides in the United States: interrupted time-series analysis. *BMJ*. 2021;375:e067726 (<https://doi.org/10.1136/bmj-2021-067726>, accessed 20 February 2025).
19. Preventing suicide: a resource for media professionals, update 2023. Geneva: World Health Organization; 2023 (<https://iris.who.int/handle/10665/372691>, accessed 20 February 2025). Licence: CC BY-NC-SA 3.0 IGO.
20. Wasserman D, Hoven CW, Wasserman C, Wall M, Eisenberg R, Hadlaczky G et al. School-based suicide prevention programmes: the SEYLE cluster-randomised, controlled trial. *Lancet*. 2015;385(9977):1536–44 ([https://doi.org/10.1016/S0140-6736\(14\)61213-7](https://doi.org/10.1016/S0140-6736(14)61213-7), accessed 20 February 2025).
21. Guidelines on mental health promotive and preventive interventions for adolescents. Geneva: World Health Organization; 2020 (<https://www.who.int/publications/i/item/9789240011854>, accessed 20 February 2025).
22. Hawton K, Witt KG, Salisbury TLT, et al. Psychosocial interventions following self-harm in adults: a systematic review and meta-analysis. *The Lancet Psychiatry* 2016; 3(8): 740-50 ([https://doi.org/10.1016/S0140-6736\(14\)61213-7](https://doi.org/10.1016/S0140-6736(14)61213-7), accessed 20 February 2025).
23. mhGAP intervention guide for mental, neurological and substance use disorders in non-specialized health settings: Mental Health Gap Action Programme (mhGAP), version 2.0. Geneva: World Health Organization; 2016 (<https://iris.who.int/handle/10665/250239>, accessed 20 February 2025).

Annex 1

Table A1.1 Number and rates of suicide, 2021

WHO African Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Algeria	Very low	UMI	Both sexes	976	2.2	2.3
			Females	292	1.3	1.4
			Males	684	3.0	3.2
Angola	Very low	LMI	Both sexes	2802	8.1	14.6
			Females	672	3.9	6.3
			Males	2130	12.5	24.4
Benin	Very low	LMI	Both sexes	956	7.1	11.8
			Females	207	3.1	5.1
			Males	749	11.1	19.3
Botswana	Very low	UMI	Both sexes	201	8.4	10.4
			Females	66	5.5	6.8
			Males	134	11.2	14.2
Burkina Faso	Very low	LI	Both sexes	1869	8.5	15.4
			Females	443	4.0	6.7
			Males	1426	13.0	26.2
Burundi	Very low	LI	Both sexes	992	7.7	14.8
			Females	254	3.9	7.0
			Males	738	11.5	24.0
Cabo Verde	Low	LMI	Both sexes	77	14.9	16.6
			Females	11	4.2	4.4
			Males	66	25.3	30.3
Cameroon	Very low	LMI	Both sexes	2287	8.5	13.8
			Females	477	3.5	6.0
			Males	1809	13.5	22.4

WHO African Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Central African Republic	Very low	LI	Both sexes	470	9.2	18.7
			Females	104	4.0	7.5
			Males	366	14.7	33.1
Chad	Very low	LI	Both sexes	929	5.2	10.7
			Females	269	3.0	6.4
			Males	660	7.4	15.4
Comoros	Very low	LMI	Both sexes	49	6.0	8.9
			Females	18	4.5	6.4
			Males	31	7.5	11.5
Congo	Very low	LMI	Both sexes	396	6.7	11.2
			Females	98	3.3	5.5
			Males	298	10.1	17.5
Côte d'Ivoire	Very low	LMI	Both sexes	2191	7.4	12.7
			Females	376	2.6	4.4
			Males	1815	12.0	20.8
Democratic Republic of the Congo	Very low	LI	Both sexes	8695	8.8	15.8
			Females	1953	3.9	6.8
			Males	6742	13.7	25.8
Equatorial Guinea	Very low	UMI	Both sexes	124	7.0	10.5
			Females	29	3.5	5.0
			Males	95	10.2	15.9
Eritrea	Very low	LI	Both sexes	443	13.2	20.2
			Females	122	7.2	10.5
			Males	321	19.4	31.7
Eswatini	Very low	LMI	Both sexes	329	27.2	31.8
			Females	62	10.1	11.4
			Males	266	45.0	54.2
Ethiopia	Very low	LI	Both sexes	7412	6.1	10.7
			Females	1864	3.1	5.1
			Males	5547	9.1	17.1

WHO African Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Gabon	Very low	UMI	Both sexes	173	7.3	10.5
			Females	28	2.4	3.5
			Males	145	12.0	17.6
Gambia	Very low	LI	Both sexes	124	4.8	8.3
			Females	40	3.1	5.4
			Males	84	6.5	11.4
Ghana	Very low	LMI	Both sexes	1729	5.3	8.1
			Females	381	2.3	3.4
			Males	1348	8.3	13.5
Guinea	Very low	LMI	Both sexes	661	4.8	8.2
			Females	246	3.5	5.7
			Males	415	6.1	11.3
Guinea-Bissau	Very low	LI	Both sexes	158	7.7	12.2
			Females	40	3.9	6.0
			Males	118	11.6	19.8
Kenya	Very low	LMI	Both sexes	2438	4.6	8.9
			Females	666	2.5	4.6
			Males	1772	6.7	14.2
Lesotho	Very low	LMI	Both sexes	648	28.7	36.7
			Females	193	16.6	20.3
			Males	455	41.4	55.8
Liberia	Very low	LI	Both sexes	314	6.0	9.6
			Females	103	3.9	6.3
			Males	210	8.0	13.3
Madagascar	Very low	LI	Both sexes	1808	6.1	10.0
			Females	515	3.5	5.5
			Males	1294	8.7	15.0
Malawi	Very low	LI	Both sexes	1517	7.6	14.2
			Females	299	2.9	5.3
			Males	1219	12.5	25.4
Mali	Very low	LI	Both sexes	931	4.2	8.3
			Females	309	2.8	5.4
			Males	621	5.5	11.4

WHO African Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Mauritania	Very low	LMI	Both sexes	128	2.7	4.9
			Females	42	1.7	3.1
			Males	87	3.7	6.9
Mauritius	High	UMI	Both sexes	135	10.5	9.6
			Females	19	3.0	2.9
			Males	115	18.0	15.9
Mozambique	Very low	LI	Both sexes	3363	10.6	19.0
			Females	974	6.0	8.8
			Males	2389	15.6	33.1
Namibia	Very low	UMI	Both sexes	241	8.6	11.1
			Females	51	3.6	4.6
			Males	190	13.9	18.4
Niger	Very low	LI	Both sexes	1097	4.5	8.9
			Females	357	3.0	5.8
			Males	740	5.9	12.1
Nigeria	Very low	LMI	Both sexes	10 912	5.0	8.8
			Females	1658	1.5	2.9
			Males	9254	8.4	15.1
Rwanda	Very low	LI	Both sexes	1159	8.7	13.6
			Females	327	4.8	7.0
			Males	832	12.8	22.4
Sao Tome and Principe	Very low	LMI	Both sexes	2	0.9	1.3
			Females	0	0.4	0.7
			Males	1	1.3	1.9
Senegal	Very low	LMI	Both sexes	1164	6.8	11.6
			Females	252	3.0	4.9
			Males	912	10.4	18.4
Seychelles	Medium	HI	Both sexes	5	4.3	3.9
			Females	1	1.7	1.7
			Males	4	6.3	6.0
Sierra Leone	Very low	LI	Both sexes	497	6.1	9.7
			Females	182	4.5	6.9
			Males	315	7.8	12.9

WHO African Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
South Africa	Medium	UMI	Both sexes	13 712	22.3	21.1
			Females	3145	9.9	9.7
			Males	10 567	35.4	32.9
South Sudan	Very low	LI	Both sexes	857	7.9	13.4
			Females	213	3.9	6.2
			Males	645	12.1	22.5
Togo	Very low	LI	Both sexes	829	9.3	15.2
			Females	175	4.0	6.4
			Males	654	14.6	24.4
Uganda	Very low	LI	Both sexes	2524	5.5	11.6
			Females	525	2.3	4.3
			Males	1999	8.8	20.9
United Republic of Tanzania	Very low	LMI	Both sexes	3207	5.1	9.1
			Females	868	2.7	4.6
			Males	2339	7.5	14.6
Zambia	Very low	LMI	Both sexes	1397	7.1	13.3
			Females	305	3.1	5.5
			Males	1091	11.2	22.6
Zimbabwe	Very low	LMI	Both sexes	2740	17.3	25.4
			Females	971	11.7	15.8
			Males	1769	23.6	38.2

WHO Region of the Americas						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Antigua and Barbuda	Medium	HI	Both sexes	1	1.5	1.4
			Females	1	1.3	1.0
			Males	1	1.6	1.6
Argentina	Medium	UMI	Both sexes	3594	7.9	7.7
			Females	770	3.4	3.4
			Males	2823	12.6	12.2
Bahamas	Medium	HI	Both sexes	13	3.3	2.9
			Females	2	0.8	0.7
			Males	11	5.9	5.4
Barbados	Low	HI	Both sexes	10	3.5	2.8
			Females	2	1.0	0.9
			Males	8	6.2	5.1
Belize	High	UMI	Both sexes	17	4.2	4.6
			Females	3	1.5	1.6
			Males	14	6.8	7.7
Bolivia	Very low	LMI	Both sexes	501	4.2	5.1
			Females	140	2.4	2.6
			Males	361	6.0	7.8
Brazil	High	UMI	Both sexes	15 907	7.6	6.9
			Females	4141	3.9	3.6
			Males	11 766	11.4	10.6
Canada	High	HI	Both sexes	3631	9.4	8.1
			Females	919	4.7	4.1
			Males	2712	14.2	12.1
Chile	High	HI	Both sexes	1496	7.7	6.8
			Females	306	3.1	2.9
			Males	1189	12.3	10.8
Colombia	High	UMI	Both sexes	2534	5.0	4.7
			Females	540	2.1	2.0
			Males	1994	7.9	7.7
Costa Rica	High	UMI	Both sexes	408	8.1	7.6
			Females	71	2.8	2.7
			Males	337	13.5	12.5

WHO Region of the Americas						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Cuba	High	UMI	Both sexes	1534	13.8	9.2
			Females	243	4.3	2.8
			Males	1291	23.5	16.2
Dominican Republic	Low	UMI	Both sexes	467	4.2	4.3
			Females	83	1.5	1.5
			Males	384	6.9	7.4
Ecuador	Medium	UMI	Both sexes	1336	7.6	7.4
			Females	233	2.6	2.6
			Males	1102	12.5	12.4
El Salvador	Low	UMI	Both sexes	478	7.6	7.8
			Females	103	3.2	3.1
			Males	374	12.6	14.0
Grenada	Medium	UMI	Both sexes	2	1.3	1.2
			Females	0	0.7	0.5
			Males	1	1.9	1.7
Guatemala	High	UMI	Both sexes	869	4.9	5.3
			Females	249	2.8	2.8
			Males	620	7.1	7.9
Guyana	Medium	HI	Both sexes	202	24.8	26.3
			Females	43	10.4	10.6
			Males	159	39.9	44.0
Haiti	Very low	LMI	Both sexes	889	7.8	9.3
			Females	291	5.1	5.5
			Males	598	10.6	13.6
Honduras	Very low	LMI	Both sexes	306	3.0	3.7
			Females	49	0.9	1.1
			Males	257	5.0	6.6
Jamaica	High	UMI	Both sexes	48	1.7	1.7
			Females	6	0.4	0.4
			Males	42	3.0	3.1
Mexico	High	UMI	Both sexes	8872	7.0	6.9
			Females	1706	2.6	2.6
			Males	7165	11.6	11.5

WHO Region of the Americas						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Nicaragua	High	LMI	Both sexes	282	4.2	4.2
			Females	57	1.7	1.6
			Males	225	6.9	6.9
Panama	High	HI	Both sexes	143	3.3	3.3
			Females	21	1.0	1.0
			Males	121	5.6	5.6
Paraguay	Medium	UMI	Both sexes	424	6.4	6.5
			Females	156	4.7	4.7
			Males	268	8.0	8.5
Peru	Medium	UMI	Both sexes	510	1.5	1.5
			Females	134	0.8	0.8
			Males	376	2.3	2.3
Saint Lucia	High	UMI	Both sexes	9	5.3	4.7
			Females	1	1.3	1.1
			Males	8	9.3	8.6
Saint Vincent and the Grenadines	High	UMI	Both sexes	0	0.4	0.3
			Females	0	0.0	0.0
			Males	0	0.8	0.6
Suriname	Medium	UMI	Both sexes	138	22.3	22.5
			Females	43	14.0	14.2
			Males	94	30.5	31.4
Trinidad and Tobago	Low	HI	Both sexes	199	13.3	11.6
			Females	35	4.6	4.1
			Males	164	22.2	19.6
United States of America	High	HI	Both sexes	53 162	15.6	14.2
			Females	10 938	6.5	5.9
			Males	42 224	24.7	22.3
Uruguay	Medium	HI	Both sexes	841	24.8	22.0
			Females	184	10.5	8.9
			Males	657	39.9	36.0

WHO Region of the Americas						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Venezuela (Bolivarian Republic of)	High	UMI	Both sexes	2437	8.6	8.9
			Females	288	2.0	2.0
			Males	2148	15.4	16.2

WHO Eastern Mediterranean Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Afghanistan	Very low	LI	Both sexes	1438	3.6	5.3
			Females	468	2.4	3.7
			Males	970	4.8	7.0
Bahrain	Low	HI	Both sexes	70	4.7	4.2
			Females	8	1.5	1.5
			Males	62	6.6	5.9
Djibouti	Very low	LMI	Both sexes	89	8.0	10.1
			Females	36	6.4	7.7
			Males	53	9.6	13.0
Egypt	Low	LMI	Both sexes	699	0.6	0.7
			Females	184	0.3	0.4
			Males	515	0.9	1.1
Iran (Islamic Republic of)	Medium	UMI	Both sexes	3603	4.1	4.1
			Females	1007	2.3	2.4
			Males	2596	5.8	5.7
Iraq	Low	UMI	Both sexes	1281	3.0	3.8
			Females	327	1.5	1.8
			Males	954	4.4	6.1

WHO Eastern Mediterranean Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Jordan	Medium	LMI	Both sexes	67	0.6	0.7
			Females	15	0.3	0.3
			Males	52	0.9	1.0
Kuwait	Low	HI	Both sexes	101	2.3	2.2
			Females	6	0.3	0.4
			Males	95	3.6	3.3
Lebanon	Low	LMI	Both sexes	40	0.7	0.7
			Females	16	0.6	0.6
			Males	24	0.9	0.9
Libya	Very low	UMI	Both sexes	357	5.0	5.1
			Females	95	2.7	2.8
			Males	262	7.2	7.4
Morocco	Very low	LMI	Both sexes	1076	2.9	3.0
			Females	444	2.4	2.4
			Males	633	3.4	3.5
Oman	Very low	HI	Both sexes	41	0.9	0.9
			Females	4	0.2	0.3
			Males	37	1.4	1.4
Pakistan	Very low	LMI	Both sexes	13 506	5.6	6.5
			Females	3711	3.2	3.4
			Males	9794	8.0	9.5
Qatar	Very low	HI	Both sexes	132	4.7	4.2
			Females	6	0.8	0.9
			Males	126	6.2	5.6
Saudi Arabia	High	HI	Both sexes	296	0.9	1.6
			Females	79	0.7	1.2
			Males	217	1.1	2.0
Somalia	Very low	LI	Both sexes	1375	8.0	15.3
			Females	525	6.1	11.4
			Males	850	9.8	19.6
Sudan	Very low	LI	Both sexes	1519	3.2	3.9
			Females	586	2.4	2.8
			Males	933	3.9	5.1

WHO Eastern Mediterranean Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Syrian Arab Republic	Low	LI	Both sexes	127	0.6	0.7
			Females	28	0.3	0.3
			Males	98	0.9	1.1
Tunisia	Low	LMI	Both sexes	222	1.8	1.8
			Females	74	1.2	1.2
			Males	148	2.5	2.5
United Arab Emirates	Low	HI	Both sexes	167	1.7	1.7
			Females	13	0.4	0.5
			Males	154	2.5	2.3
Yemen	Very low	LI	Both sexes	1564	4.2	5.2
			Females	520	2.8	3.3
			Males	1044	5.6	7.2

WHO European Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Albania	Low	UMI	Both sexes	78	2.8	2.1
			Females	33	2.3	1.7
			Males	45	3.2	2.6
Armenia	High	UMI	Both sexes	69	2.5	1.9
			Females	14	1.0	0.7
			Males	55	4.2	3.6
Austria	High	HI	Both sexes	1301	14.5	10.2
			Females	262	5.7	4.1
			Males	1040	23.6	16.7
Azerbaijan	Very low	UMI	Both sexes	160	1.6	1.6
			Females	43	0.8	0.8
			Males	117	2.3	2.4
Belarus	High	UMI	Both sexes	1444	15.6	11.7
			Females	250	5.1	3.3
			Males	1194	27.7	22.1
Belgium	High	HI	Both sexes	2124	18.4	14.3
			Females	627	10.7	8.0
			Males	1497	26.2	20.8
Bosnia and Herzegovina	Medium	UMI	Both sexes	291	9.0	5.7
			Females	80	4.7	2.4
			Males	211	13.7	9.5
Bulgaria	Medium	HI	Both sexes	654	9.5	6.3
			Females	149	4.2	2.6
			Males	505	15.2	10.4
Croatia	High	HI	Both sexes	615	15.7	10.6
			Females	141	6.9	4.2
			Males	474	25.1	17.6
Cyprus	High	HI	Both sexes	40	3.0	2.7
			Females	14	2.2	1.6
			Males	26	3.8	3.7

WHO European Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Czechia	High	HI	Both sexes	1403	13.3	10.3
			Females	303	5.7	4.6
			Males	1100	21.2	16.3
Denmark	High	HI	Both sexes	613	10.5	7.6
			Females	186	6.3	4.3
			Males	427	14.7	11.0
Estonia	High	HI	Both sexes	199	14.9	12.5
			Females	40	5.8	5.3
			Males	159	25.1	20.4
Finland	High	HI	Both sexes	807	14.6	12.6
			Females	246	8.8	7.9
			Males	562	20.5	17.4
France	Medium	HI	Both sexes	10 957	16.6	11.8
			Females	3242	9.5	6.0
			Males	7715	24.1	18.0
Georgia	Medium	UMI	Both sexes	193	5.1	4.6
			Females	26	1.3	1.0
			Males	167	9.5	8.9
Germany	High	HI	Both sexes	10 799	12.9	8.2
			Females	3213	7.6	4.2
			Males	7587	18.4	12.2
Greece	Medium	HI	Both sexes	495	4.7	3.4
			Females	89	1.6	1.2
			Males	407	7.9	5.8
Hungary	High	HI	Both sexes	1597	16.5	11.4
			Females	378	7.5	4.8
			Males	1219	26.2	19.0
Iceland	High	HI	Both sexes	44	11.9	10.8
			Females	17	9.3	8.4
			Males	27	14.4	13.0
Ireland	High	HI	Both sexes	430	8.5	7.7
			Females	116	4.6	4.2
			Males	313	12.6	11.3

WHO European Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Israel	High	HI	Both sexes	391	4.4	4.4
			Females	110	2.4	2.3
			Males	280	6.3	6.5
Italy	High	HI	Both sexes	4176	7.0	4.6
			Females	1108	3.6	2.2
			Males	3068	10.5	7.1
Kazakhstan	High	UMI	Both sexes	2882	14.6	15.2
			Females	540	5.3	5.5
			Males	2343	24.4	26.2
Kyrgyzstan	High	LMI	Both sexes	464	6.8	7.8
			Females	108	3.1	3.6
			Males	355	10.5	12.4
Latvia	High	HI	Both sexes	287	15.2	12.0
			Females	47	4.6	3.0
			Males	240	27.5	22.1
Lithuania	High	HI	Both sexes	618	22.1	16.8
			Females	136	9.2	5.9
			Males	483	36.7	28.8
Luxembourg	High	HI	Both sexes	53	8.2	6.2
			Females	14	4.3	2.9
			Males	39	12.1	9.6
Malta	High	HI	Both sexes	34	6.5	5.6
			Females	11	4.4	3.3
			Males	23	8.4	7.5
Montenegro	Low	UMI	Both sexes	68	11.3	8.9
			Females	19	6.0	4.3
			Males	50	17.1	14.1
Netherlands (Kingdom of the)	High	HI	Both sexes	2040	11.5	9.1
			Females	680	7.6	5.9
			Males	1359	15.4	12.5
North Macedonia	Medium	UMI	Both sexes	107	5.8	4.6
			Females	36	3.8	3.0
			Males	71	7.9	6.4

WHO European Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Norway	High	HI	Both sexes	713	13.2	11.4
			Females	199	7.4	6.3
			Males	515	18.9	16.3
Poland	Medium	HI	Both sexes	5199	13.7	11.5
			Females	753	3.8	3.2
			Males	4446	24.1	20.2
Portugal	High	HI	Both sexes	1197	11.5	7.2
			Females	323	5.9	3.5
			Males	874	17.7	11.4
Republic of Moldova	High	UMI	Both sexes	428	14.1	11.9
			Females	82	5.0	4.0
			Males	346	24.8	22.1
Romania	High	HI	Both sexes	1852	9.6	7.2
			Females	327	3.3	2.3
			Males	1526	16.4	12.5
Russian Federation	Medium	HI	Both sexes	31 172	21.4	17.9
			Females	6320	8.1	6.1
			Males	24 852	36.7	31.6
Serbia	Medium	UMI	Both sexes	1035	15.1	9.4
			Females	287	8.0	4.4
			Males	748	23.0	15.0
Slovakia	High	HI	Both sexes	539	9.9	7.6
			Females	84	3.0	2.3
			Males	455	17.1	13.5
Slovenia	High	HI	Both sexes	398	18.9	13.2
			Females	81	7.7	5.1
			Males	318	29.9	21.4
Spain	High	HI	Both sexes	4160	8.7	6.2
			Females	1085	4.5	3.2
			Males	3075	13.1	9.4
Sweden	High	HI	Both sexes	1442	13.8	11.7
			Females	450	8.7	7.1
			Males	992	18.9	16.2

WHO European Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Switzerland	High	HI	Both sexes	1221	14.0	9.9
			Females	403	9.2	6.0
			Males	818	18.9	13.9
Tajikistan	Low	LMI	Both sexes	242	2.4	3.1
			Females	68	1.3	1.7
			Males	174	3.6	4.5
Türkiye	Medium	UMI	Both sexes	2332	2.7	2.6
			Females	473	1.1	1.0
			Males	1859	4.3	4.2
Turkmenistan	Very low	UMI	Both sexes	484	6.8	7.5
			Females	126	3.5	3.9
			Males	358	10.3	11.7
Ukraine	Medium	UMI	Both sexes	9397	21.2	17.0
			Females	1625	6.9	4.7
			Males	7773	37.7	30.9
United Kingdom of Great Britain and Northern Ireland	High	HI	Both sexes	6461	9.5	8.8
			Females	1581	4.6	4.1
			Males	4880	14.7	13.5
Uzbekistan	Medium	LMI	Both sexes	2857	8.3	9.1
			Females	1029	6.1	6.5
			Males	1828	10.6	11.8

WHO South-East Asia Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Bangladesh	Very low	LMI	Both sexes	4714	2.8	2.9
			Females	1701	2.0	1.9
			Males	3013	3.6	3.9
Bhutan	Very low	LMI	Both sexes	38	4.9	5.1
			Females	12	3.4	3.6
			Males	26	6.2	6.5
Democratic People's Republic of Korea	Very low	LI	Both sexes	2442	9.3	7.6
			Females	1078	8.1	6.3
			Males	1363	10.5	9.1
India	Very low	LMI	Both sexes	177 567	12.6	12.6
			Females	76 848	11.2	11.1
			Males	100 719	13.8	14.1
Indonesia	Very low	UMI	Both sexes	3333	1.2	1.3
			Females	1257	0.9	0.9
			Males	2076	1.5	1.7
Maldives	Low	UMI	Both sexes	8	1.5	2.0
			Females	1	0.3	0.4
			Males	7	2.2	3.2
Myanmar	Very low	LMI	Both sexes	1545	2.9	2.9
			Females	274	1.0	1.0
			Males	1271	4.8	4.9
Nepal	Very low	LMI	Both sexes	2989	10.1	11.1
			Females	1091	7.2	7.2
			Males	1898	13.3	15.8
Sri Lanka	Medium	LMI	Both sexes	3347	14.7	14.0
			Females	685	5.9	5.6
			Males	2661	24.2	23.6
Thailand	Low	UMI	Both sexes	11 899	16.6	15.4
			Females	2011	5.5	4.8
			Males	9887	28.2	26.2

Timor-Leste	Very low	LMI	Both sexes	48	3.6	4.2
			Females	15	2.3	2.6
			Males	33	4.8	5.8

WHO Western Pacific Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Australia	High	HI	Both sexes	3394	13.1	11.7
			Females	880	6.7	6.1
			Males	2514	19.5	17.5
Brunei Darussalam	High	HI	Both sexes	14	3.0	2.8
			Females	1	0.6	0.7
			Males	12	5.2	4.6
Cambodia	Very low	LMI	Both sexes	764	4.5	5.0
			Females	237	2.7	2.9
			Males	526	6.3	7.4
China	Very low	UMI	Both sexes	127 601	8.9	6.8
			Females	52 479	7.5	5.3
			Males	75 122	10.3	8.5
Fiji	Low	UMI	Both sexes	76	8.3	8.5
			Females	25	5.3	5.4
			Males	51	11.2	11.8
Japan	High	HI	Both sexes	21 904	17.4	14.7
			Females	7417	11.5	9.8
			Males	14 487	23.6	19.6
Kiribati	Very low	LMI	Both sexes	22	17.5	18.7
			Females	3	4.8	5.2
			Males	19	31.0	33.9
Lao People's Democrati	Very low	LMI	Both sexes	341	4.6	5.0
			Females	96	2.6	2.7
			Males	245	6.5	7.4

WHO Western Pacific Region						
Country	Data quality ^a	Income group ^b	Sex	Number of suicides, ^c 2021	Crude suicide rate, ^d 2021	Age-standardized suicide rate, ^d 2021
Malaysia	Low	UMI	Both sexes	1938	5.7	5.6
			Females	355	2.2	2.2
			Males	1584	8.8	8.8
Micronesia (Federated States of)	Low	LMI	Both sexes	22	19.8	20.8
			Females	5	8.6	8.9
			Males	17	31.1	33.1
Mongolia	High	UMI	Both sexes	618	18.5	19.8
			Females	81	4.8	5.4
			Males	537	32.2	34.5
New Zealand	High	HI	Both sexes	609	11.9	11.1
			Females	156	6.1	5.8
			Males	453	17.9	16.5
Papua New Guinea	Very low	LMI	Both sexes	182	1.8	2.1
			Females	68	1.4	1.6
			Males	114	2.2	2.7
Philippines	Medium	LMI	Both sexes	3947	3.5	4.0
			Females	1000	1.8	2.1
			Males	2946	5.2	6.1
Republic of Korea	High	HI	Both sexes	14 273	27.5	20.6
			Females	4399	16.9	13.6
			Males	9874	38.2	28.2
Samoa	Very low	LMI	Both sexes	26	12.0	14.3
			Females	7	7.0	8.0
			Males	18	17.0	20.8
Singapore	High	HI	Both sexes	449	8.1	7.0
			Females	131	4.9	4.3
			Males	318	11.1	9.6
Solomon Islands	Low	LMI	Both sexes	149	19.6	22.5
			Females	36	9.6	10.9
			Males	114	29.2	33.7

Tonga	Very low	UMI	Both sexes	5	4.7	5.4
			Females	2	3.5	3.8
			Males	3	6.0	7.3
Vanuatu	Very low	LMI	Both sexes	48	15.7	18.2
			Females	10	6.6	7.6
			Males	38	24.7	28.9
Viet Nam	Very low	LMI	Both sexes	7222	7.3	7.0
			Females	2191	4.3	3.9
			Males	5031	10.4	10.5

^a High quality = Multiple years of national death registration data with high completeness and quality of cause-of-death assignment are available. Estimates for these countries may be compared and time series may be used for priority setting and policy evaluation. Medium and low quality = Multiple years of death registration data are available. Data have low completeness and/or issues with cause-of-death assignment which likely affect estimated deaths by cause and time trends. Estimates may be used for priority setting. Use estimates for programme evaluation with caution, as improvements in the vital registration system may affect the estimated trends in cause-specific mortality. Comparisons among countries should be interpreted with caution. Medium quality denotes moderate quality issues and low quality denotes severe quality issues. Low quality = Death registration data are unavailable or unusable due to quality issues. Estimates of mortality by cause should be interpreted with caution. Estimates may be used for priority setting, however, they are not likely to be informative for policy evaluation or comparisons among countries.

^b World Bank income groups, 2024. HI = high-income, LI = low-income, LMI = lower-middle-income, UMI = upper-middle-income

^c All ages

^d All ages, per 100 000 population

Annex 2

Table A.2.1 Suicide rate by age groups, 2021

WHO African Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Algeria	Both sexes	3.3	3.4	2.1	3.7
	Females	1.8	2.1	1.2	2.9
	Males	4.6	4.7	2.9	4.7
Angola	Both sexes	6.1	13.3	32.6	65.4
	Females	3.5	5.8	14.1	25.6
	Males	8.7	21.1	53.6	120.7
Benin	Both sexes	5.5	11.4	25.5	48.9
	Females	2.6	3.4	12.5	24.6
	Males	8.4	19.6	39.8	81.2
Botswana	Both sexes	7.3	15.2	19.3	21.9
	Females	3.9	11.3	12.1	14.7
	Males	10.7	19.0	27.0	32.4
Burkina Faso	Both sexes	6.8	13.6	34.1	68.5
	Females	4.2	4.6	15.0	32.0
	Males	9.4	22.9	57.3	125.0
Burundi	Both sexes	6.2	12.0	32.7	70.2
	Females	4.2	4.3	15.8	34.5
	Males	8.2	19.9	51.4	121.0
Cabo Verde	Both sexes	8.5	18.3	39.0	52.1
	Females	1.7	3.6	11.1	20.7
	Males	14.6	31.6	69.3	109.4
Cameroon	Both sexes	7.2	14.4	28.5	51.7
	Females	2.9	4.7	14.2	26.6
	Males	11.5	24.2	43.7	84.6

WHO African Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Central African Republic	Both sexes	8.2	15.3	47.7	72.3
	Females	3.2	5.0	20.3	31.1
	Males	13.4	28.3	85.1	122.4
Chad	Both sexes	3.9	8.8	24.8	49.4
	Females	2.3	3.3	17.3	33.8
	Males	5.5	14.3	33.0	69.0
Comoros	Both sexes	4.2	7.0	19.3	46.5
	Females	4.0	4.0	14.9	33.5
	Males	4.4	10.0	23.6	61.8
Congo	Both sexes	5.9	9.8	22.6	55.0
	Females	2.9	4.4	11.5	28.0
	Males	8.8	15.1	34.4	88.8
Côte d'Ivoire	Both sexes	5.7	11.5	26.7	58.1
	Females	2.6	3.3	9.9	19.2
	Males	8.6	19.3	41.8	99.9
Democratic Republic of the Congo	Both sexes	6.1	13.5	37.0	69.4
	Females	2.7	4.0	18.6	31.3
	Males	9.6	23.2	57.1	117.6
Equatorial Guinea	Both sexes	5.5	8.9	21.1	52.3
	Females	3.5	3.6	10.9	22.2
	Males	7.3	12.8	30.2	88.3
Eritrea	Both sexes	10.1	19.4	43.3	87.1
	Females	5.0	8.8	24.5	47.9
	Males	15.2	30.8	66.3	140.4
Eswatini	Both sexes	29.1	55.7	44.4	32.3
	Females	14.0	15.6	16.0	19.2
	Males	43.9	96.5	81.9	53.1
Ethiopia	Both sexes	4.4	8.3	22.1	60.7
	Females	2.6	2.9	11.4	30.1
	Males	6.1	13.8	34.1	100.1

WHO African Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Gabon	Both sexes	4.7	9.7	21.8	50.8
	Females	1.5	2.4	7.8	19.8
	Males	7.9	16.2	35.4	88.6
Gambia	Both sexes	4.2	6.2	17.8	42.4
	Females	2.4	2.6	13.4	31.7
	Males	6.0	9.9	22.9	55.1
Ghana	Both sexes	3.6	6.9	17.0	41.0
	Females	1.6	2.9	7.2	16.3
	Males	5.6	10.9	27.4	74.1
Guinea	Both sexes	3.6	6.6	19.1	38.6
	Females	2.7	3.6	14.5	27.8
	Males	4.4	9.9	24.8	53.9
Guinea-Bissau	Both sexes	6.3	13.2	25.2	46.0
	Females	3.3	5.1	14.1	24.1
	Males	9.3	21.7	40.2	78.3
Kenya	Both sexes	2.5	4.9	19.8	60.9
	Females	1.1	3.0	9.7	32.7
	Males	4.0	6.7	30.6	101.2
Lesotho	Both sexes	24.3	62.1	61.2	45.4
	Females	14.7	34.6	30.9	31.9
	Males	34.1	91.2	101.5	72.3
Liberia	Both sexes	4.7	7.7	21.2	44.6
	Females	3.5	3.4	15.7	31.5
	Males	5.9	12.0	27.4	59.9
Madagascar	Both sexes	4.7	7.9	21.8	49.9
	Females	3.1	3.5	12.7	27.2
	Males	6.3	12.3	31.2	77.4
Malawi	Both sexes	5.9	12.2	31.8	68.2
	Females	2.3	3.2	12.7	29.9
	Males	9.5	22.0	55.9	127.7
Mali	Both sexes	3.2	6.9	18.9	38.9
	Females	2.8	3.1	13.8	26.0
	Males	3.6	10.6	24.3	54.8

WHO African Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Mauritania	Both sexes	2.0	3.2	10.8	28.3
	Females	0.9	1.7	8.1	18.1
	Males	3.0	4.8	13.9	40.4
Mauritius	Both sexes	16.0	13.6	10.1	3.9
	Females	5.2	2.8	3.4	1.6
	Males	26.2	23.7	17.0	6.8
Mozambique	Both sexes	11.0	18.5	36.5	81.5
	Females	8.3	8.9	14.1	34.0
	Males	13.6	29.5	70.2	160.6
Namibia	Both sexes	8.3	17.0	19.2	19.5
	Females	2.8	6.5	8.9	11.8
	Males	13.9	27.9	32.1	33.8
Niger	Both sexes	3.0	6.7	20.9	42.5
	Females	2.1	2.9	15.7	29.9
	Males	3.8	10.3	26.0	58.8
Nigeria	Both sexes	3.0	8.2	17.7	45.6
	Females	0.9	1.2	7.4	17.8
	Males	4.9	15.1	28.4	76.8
Rwanda	Both sexes	6.2	11.1	29.2	66.4
	Females	3.6	4.9	15.5	35.1
	Males	8.8	17.8	46.3	117.1
Sao Tome and Principe	Both sexes	0.8	1.3	2.7	4.4
	Females	0.2	0.7	1.7	2.4
	Males	1.3	1.9	3.7	7.2
Senegal	Both sexes	4.5	7.9	25.0	69.3
	Females	2.3	2.8	11.4	29.4
	Males	6.7	13.1	38.9	110.0
Seychelles	Both sexes	3.8	5.2	5.7	12.0
	Females	4.0	1.5	1.2	3.0
	Males	3.7	7.6	9.9	25.1
Sierra Leone	Both sexes	4.6	8.5	21.5	41.0
	Females	3.9	5.1	16.4	29.3
	Males	5.4	12.0	27.2	56.1

WHO African Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
South Africa	Both sexes	26.5	40.7	20.1	15.2
	Females	8.1	17.5	13.9	14.7
	Males	45.0	64.4	27.8	16.1
South Sudan	Both sexes	4.9	10.6	31.9	66.8
	Females	3.0	3.6	14.6	34.1
	Males	6.7	18.5	52.6	116.1
Togo	Both sexes	6.7	14.5	33.9	63.0
	Females	3.3	4.8	15.2	29.1
	Males	10.0	24.0	53.5	102.1
Uganda	Both sexes	4.3	9.3	25.7	60.6
	Females	2.1	3.1	9.0	24.5
	Males	6.5	15.6	45.1	118.5
United Republic of Tanzania	Both sexes	4.0	8.4	18.9	43.1
	Females	2.3	3.7	9.9	23.2
	Males	5.6	13.2	29.7	71.2
Zambia	Both sexes	6.1	13.4	28.8	52.8
	Females	3.1	4.9	11.6	25.2
	Males	9.1	22.0	48.9	97.7
Zimbabwe	Both sexes	16.7	37.3	46.7	56.0
	Females	11.6	24.0	25.6	36.2
	Males	22.3	54.4	75.3	85.6

WHO Region of the Americas					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Antigua and Barbuda	Both sexes	2.8	0.6	0.0	10.5
	Females	0.0	0.0	0.0	17.1
	Males	5.5	1.2	0.0	0.0
Argentina	Both sexes	13.2	8.7	7.4	12.0
	Females	6.2	3.7	2.6	3.2
	Males	19.9	13.6	12.5	25.1
Bahamas	Both sexes	3.4	4.6	3.7	4.7
	Females	0.9	1.2	0.9	1.0
	Males	6.0	8.3	6.8	10.0
Barbados	Both sexes	3.3	3.1	3.9	10.4
	Females	1.4	0.7	1.1	2.5
	Males	5.1	5.7	7.0	21.5
Belize	Both sexes	3.5	6.9	6.9	12.9
	Females	1.3	2.2	3.2	2.6
	Males	5.7	11.7	10.7	23.7
Bolivia	Both sexes	2.3	4.0	15.4	14.2
	Females	2.4	2.3	6.3	4.8
	Males	2.3	5.6	25.0	26.3
Brazil	Both sexes	7.3	8.8	10.8	16.0
	Females	4.1	4.3	5.1	7.3
	Males	10.4	13.4	17.1	27.5
Canada	Both sexes	9.1	11.3	12.9	9.9
	Females	4.6	5.2	6.4	5.5
	Males	13.4	17.4	19.7	15.3
Chile	Both sexes	8.6	9.7	8.6	11.7
	Females	4.3	4.2	2.6	3.7
	Males	12.7	15.1	14.8	21.9
Colombia	Both sexes	6.4	4.5	6.5	10.9
	Females	3.2	1.8	1.8	3.1
	Males	9.6	7.3	11.7	21.3
Costa Rica	Both sexes	9.4	12.3	8.7	5.6
	Females	3.8	4.4	2.0	0.6
	Males	14.8	20.2	16.2	12.0

WHO Region of the Americas					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Cuba	Both sexes	5.2	8.4	21.5	41.0
	Females	2.0	1.8	7.8	10.2
	Males	8.3	14.8	36.4	79.8
Dominican Republic	Both sexes	4.2	5.3	7.7	10.9
	Females	1.6	1.8	2.4	3.8
	Males	6.7	8.7	13.5	20.8
Ecuador	Both sexes	12.2	9.0	7.2	8.1
	Females	4.9	2.1	2.1	1.3
	Males	19.2	15.9	12.5	17.0
El Salvador	Both sexes	7.9	12.6	10.2	9.6
	Females	4.2	3.5	3.2	3.4
	Males	11.4	23.8	20.2	19.1
Grenada	Both sexes	0.0	2.4	1.4	4.5
	Females	0.0	0.0	0.0	7.8
	Males	0.0	4.7	2.7	0.0
Guatemala	Both sexes	7.9	6.7	5.9	6.6
	Females	5.1	3.3	2.5	2.0
	Males	10.8	10.3	9.9	12.0
Guyana	Both sexes	26.1	37.0	44.8	36.6
	Females	16.0	10.2	16.3	12.0
	Males	36.2	66.9	77.1	71.3
Haiti	Both sexes	6.6	13.5	16.7	19.3
	Females	7.4	6.6	8.1	7.6
	Males	5.7	20.5	26.2	35.9
Honduras	Both sexes	2.9	4.1	7.1	11.3
	Females	1.3	1.3	1.7	1.7
	Males	4.4	6.9	13.1	23.7
Jamaica	Both sexes	1.2	1.7	2.6	8.3
	Females	0.3	0.4	0.7	1.6
	Males	2.1	3.0	4.8	16.9
Mexico	Both sexes	11.5	9.2	5.6	7.5
	Females	5.1	2.5	1.6	3.0
	Males	17.9	16.7	10.2	13.0

WHO Region of the Americas					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Nicaragua	Both sexes	6.7	6.1	3.4	3.1
	Females	3.3	0.9	1.2	1.6
	Males	10.2	11.6	5.9	5.3
Panama	Both sexes	3.9	3.5	5.0	7.2
	Females	1.8	0.3	0.8	2.1
	Males	6.0	6.7	9.4	13.8
Paraguay	Both sexes	9.8	6.4	9.1	7.5
	Females	7.6	5.0	4.8	3.4
	Males	11.9	7.8	13.8	12.9
Peru	Both sexes	1.6	1.8	2.6	2.3
	Females	1.3	0.7	0.9	0.8
	Males	2.0	2.8	4.3	4.1
Saint Lucia	Both sexes	5.5	6.1	6.3	12.6
	Females	1.0	1.4	2.0	2.9
	Males	9.9	10.9	10.9	26.5
Saint Vincent and the Grenadines	Both sexes	0.0	0.0	1.9	0.0
	Females	0.0	0.0	0.0	0.0
	Males	0.0	0.0	3.6	0.0
Suriname	Both sexes	36.0	26.3	25.3	35.2
	Females	18.4	20.6	16.7	14.5
	Males	53.2	31.7	34.4	67.2
Trinidad and Tobago	Both sexes	12.0	17.6	17.4	19.3
	Females	5.3	5.4	6.0	6.2
	Males	18.5	29.7	29.8	38.0
United States of America	Both sexes	18.4	19.9	17.7	19.7
	Females	7.1	8.3	8.2	6.2
	Males	28.9	30.8	27.3	36.8
Uruguay	Both sexes	28.5	28.3	31.0	39.1
	Females	8.2	12.4	15.3	16.7
	Males	48.0	44.1	48.3	77.9

WHO Region of the Americas					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Venezuela (Bolivarian Republic of)	Both sexes	15.7	9.9	9.1	11.7
	Females	2.8	2.5	2.5	2.4
	Males	27.8	17.7	16.4	25.0

WHO Eastern Mediterranean Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Afghanistan	Both sexes	4.8	7.7	8.0	10.9
	Females	3.0	4.0	8.0	8.1
	Males	6.6	11.4	8.0	14.8
Bahrain	Both sexes	6.8	5.4	4.1	10.7
	Females	2.1	1.8	1.6	3.7
	Males	9.5	7.0	6.0	17.4
Djibouti	Both sexes	5.9	9.3	19.4	47.6
	Females	5.3	7.4	14.3	32.7
	Males	6.4	11.3	25.0	66.7
Egypt	Both sexes	0.9	0.7	1.2	1.7
	Females	0.5	0.3	0.6	0.8
	Males	1.3	1.0	1.8	2.9
Iran (Islamic Republic of)	Both sexes	8.3	4.7	3.2	3.2
	Females	5.4	2.1	1.8	2.5
	Males	11.1	7.2	4.5	3.9
Iraq	Both sexes	3.7	4.8	6.8	7.2
	Females	2.0	2.6	2.5	2.8
	Males	5.3	7.0	11.9	14.2
Jordan	Both sexes	0.9	0.8	0.8	1.4
	Females	0.4	0.4	0.4	1.0
	Males	1.4	1.2	1.3	1.8

WHO Eastern Mediterranean Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Kuwait	Both sexes	3.8	2.7	2.2	3.5
	Females	0.6	0.3	0.4	1.3
	Males	6.1	4.0	3.1	5.2
Lebanon	Both sexes	1.1	0.7	0.8	1.6
	Females	1.1	0.5	0.4	1.1
	Males	1.2	0.8	1.3	2.4
Libya	Both sexes	7.0	7.8	5.7	5.1
	Females	4.0	4.0	2.9	3.3
	Males	9.8	11.4	8.6	7.3
Morocco	Both sexes	3.8	3.6	4.3	5.1
	Females	3.2	3.3	3.2	3.4
	Males	4.5	3.9	5.5	7.3
Oman	Both sexes	1.2	1.2	1.2	2.0
	Females	0.2	0.4	0.4	0.9
	Males	1.9	1.6	1.7	3.3
Pakistan	Both sexes	9.6	8.3	7.2	9.2
	Females	6.6	3.7	3.1	4.3
	Males	12.4	12.8	11.3	14.8
Qatar	Both sexes	6.9	5.1	4.2	8.9
	Females	1.3	0.8	1.2	3.5
	Males	8.9	6.4	5.3	13.5
Saudi Arabia	Both sexes	1.3	0.4	1.7	13.5
	Females	0.4	0.4	1.1	13.7
	Males	2.0	0.5	2.0	13.4
Somalia	Both sexes	6.6	14.0	35.3	64.8
	Females	5.0	8.6	28.2	52.0
	Males	8.1	19.5	43.3	81.4
Sudan	Both sexes	5.0	5.7	4.3	4.7
	Females	4.3	3.7	2.7	3.0
	Males	5.6	8.0	6.0	6.2
Syrian Arab Republic	Both sexes	0.8	0.9	0.9	1.2
	Females	0.4	0.3	0.3	0.6
	Males	1.2	1.4	1.7	1.9

WHO Eastern Mediterranean Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Tunisia	Both sexes	2.5	2.2	2.1	4.2
	Females	1.5	1.5	1.4	2.9
	Males	3.5	3.0	2.9	6.0
United Arab Emirates	Both sexes	1.7	2.1	2.3	4.8
	Females	0.4	0.4	0.6	3.1
	Males	2.5	2.8	3.2	6.2
Yemen	Both sexes	6.4	7.7	6.5	5.9
	Females	5.3	4.0	3.6	3.7
	Males	7.5	11.4	9.6	9.0

WHO European Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Albania	Both sexes	2.1	2.0	3.0	9.7
	Females	2.1	1.5	2.3	8.3
	Males	2.2	2.6	3.9	11.3
Armenia	Both sexes	1.1	1.8	3.4	11.1
	Females	0.3	0.0	0.8	6.8
	Males	1.9	3.9	7.0	19.2
Austria	Both sexes	9.0	12.8	17.1	34.1
	Females	3.2	5.1	8.0	10.4
	Males	14.5	20.4	26.4	66.7
Azerbaijan	Both sexes	2.3	1.5	2.1	3.7
	Females	0.9	0.6	1.3	3.2
	Males	3.6	2.4	3.2	4.5
Belarus	Both sexes	6.2	16.4	24.6	29.8
	Females	1.8	3.9	6.4	13.6
	Males	10.3	28.9	48.0	67.9

WHO European Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Belgium	Both sexes	10.1	22.2	26.3	28.6
	Females	5.5	11.3	16.2	17.2
	Males	14.5	33.1	36.6	43.9
Bosnia and Herzegovina	Both sexes	4.6	5.0	12.1	24.0
	Females	1.1	1.9	5.8	13.2
	Males	7.7	8.0	19.4	44.4
Bulgaria	Both sexes	4.6	7.0	13.5	20.9
	Females	1.4	3.4	5.2	9.1
	Males	7.6	10.4	22.6	40.2
Croatia	Both sexes	7.3	12.7	22.6	31.2
	Females	2.2	4.8	9.1	15.4
	Males	12.2	20.3	37.3	57.4
Cyprus	Both sexes	1.6	3.7	4.0	3.6
	Females	0.0	3.3	3.3	3.3
	Males	3.2	4.0	4.8	3.9
Czechia	Both sexes	9.8	13.9	17.2	23.7
	Females	5.2	5.2	8.4	7.7
	Males	14.0	22.0	26.3	47.7
Denmark	Both sexes	6.4	10.2	13.1	22.8
	Females	3.8	5.0	8.1	13.9
	Males	8.8	15.3	18.0	33.7
Estonia	Both sexes	17.6	16.2	16.6	23.2
	Females	12.5	4.5	3.5	10.3
	Males	22.4	27.0	32.0	50.4
Finland	Both sexes	14.1	18.5	18.3	16.6
	Females	9.1	11.2	11.0	8.9
	Males	18.8	25.4	25.8	26.9
France	Both sexes	7.2	16.3	23.6	34.8
	Females	4.1	6.7	13.2	22.2
	Males	10.2	26.2	34.8	52.4
Georgia	Both sexes	5.8	5.4	6.3	10.8
	Females	0.7	1.2	1.5	3.9
	Males	10.7	10.0	12.6	25.1

WHO European Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Germany	Both sexes	6.3	9.5	15.8	30.3
	Females	2.9	4.4	8.7	19.3
	Males	9.5	14.5	23.0	45.2
Greece	Both sexes	2.1	5.1	6.8	6.9
	Females	1.0	1.5	2.2	2.4
	Males	3.1	8.7	11.8	12.9
Hungary	Both sexes	7.9	14.0	25.0	33.4
	Females	3.5	5.6	9.9	16.3
	Males	12.0	22.0	42.3	64.0
Iceland	Both sexes	14.3	14.3	18.6	4.6
	Females	13.3	8.4	13.0	8.7
	Males	15.2	19.5	24.0	0.0
Ireland	Both sexes	7.9	11.4	13.1	8.0
	Females	5.0	5.8	6.6	4.0
	Males	10.6	17.2	19.7	12.6
Israel	Both sexes	4.3	6.4	6.9	7.9
	Females	1.5	3.5	3.5	6.7
	Males	6.9	9.3	10.5	9.4
Italy	Both sexes	3.6	6.3	8.6	13.2
	Females	1.6	2.9	4.2	7.3
	Males	5.4	9.6	13.3	21.2
Kazakhstan	Both sexes	15.1	23.1	19.7	29.7
	Females	7.7	5.6	5.7	17.8
	Males	22.2	40.9	36.9	54.9
Kyrgyzstan	Both sexes	8.5	9.6	10.6	18.3
	Females	5.2	2.7	4.4	12.0
	Males	11.9	16.6	17.7	27.8
Latvia	Both sexes	8.9	20.1	17.5	23.7
	Females	4.0	3.6	2.9	12.4
	Males	13.5	36.3	35.8	49.1
Lithuania	Both sexes	15.8	22.4	30.7	35.3
	Females	4.0	7.5	11.8	17.9
	Males	26.8	36.2	53.8	73.5

WHO European Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Luxembourg	Both sexes	3.3	9.1	13.0	16.0
	Females	0.0	4.7	7.8	8.5
	Males	6.4	13.4	18.0	25.7
Malta	Both sexes	5.8	8.0	8.9	5.5
	Females	1.0	8.0	4.2	5.2
	Males	9.9	8.0	13.5	5.8
Montenegro	Both sexes	6.9	12.1	16.1	24.4
	Females	3.8	4.8	7.7	15.5
	Males	9.8	19.6	25.4	39.6
Netherlands (Kingdom of the)	Both sexes	8.9	12.0	16.6	16.4
	Females	5.5	7.3	10.8	11.9
	Males	12.1	16.6	22.5	22.0
North Macedonia	Both sexes	4.7	5.7	8.7	9.9
	Females	1.9	4.6	6.1	4.3
	Males	7.5	6.8	11.5	18.1
Norway	Both sexes	12.4	16.6	17.5	15.8
	Females	6.3	8.8	10.0	9.7
	Males	18.0	24.0	24.8	23.2
Poland	Both sexes	12.7	16.2	19.0	14.5
	Females	3.6	3.5	5.7	5.1
	Males	21.3	28.5	33.7	30.2
Portugal	Both sexes	4.2	9.4	14.9	24.9
	Females	1.9	4.5	7.7	12.2
	Males	6.5	14.7	23.1	42.9
Republic of Moldova	Both sexes	10.5	15.6	21.5	27.1
	Females	4.1	4.9	4.7	15.3
	Males	17.2	27.1	43.9	52.0
Romania	Both sexes	5.3	9.4	15.1	15.7
	Females	1.3	2.8	5.1	6.1
	Males	9.1	15.7	26.4	31.6
Russian Federation	Both sexes	15.1	29.3	25.9	32.5
	Females	5.4	8.6	8.8	18.4
	Males	24.4	50.6	48.3	64.4

WHO European Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Serbia	Both sexes	4.1	10.8	21.4	38.6
	Females	0.8	5.6	10.1	21.8
	Males	7.4	16.2	34.3	63.0
Slovakia	Both sexes	6.1	9.8	13.0	22.5
	Females	2.0	2.3	4.4	6.2
	Males	10.0	17.0	22.4	50.0
Slovenia	Both sexes	8.4	17.5	25.1	40.5
	Females	3.0	6.9	10.7	14.3
	Males	13.2	27.0	39.5	78.6
Spain	Both sexes	4.4	9.0	11.7	15.3
	Females	2.2	4.4	6.3	7.0
	Males	6.4	13.6	17.2	26.7
Sweden	Both sexes	14.2	15.3	18.1	19.9
	Females	7.5	9.2	11.4	13.5
	Males	20.4	21.0	24.6	27.6
Switzerland	Both sexes	8.8	11.2	17.9	33.4
	Females	5.1	6.5	10.8	23.7
	Males	12.3	15.7	25.0	46.1
Tajikistan	Both sexes	2.8	3.9	5.1	7.9
	Females	1.8	1.5	2.9	6.2
	Males	3.8	6.5	7.4	10.3
Türkiye	Both sexes	3.8	2.7	3.4	5.3
	Females	1.6	1.0	1.2	2.9
	Males	6.0	4.4	5.8	8.5
Turkmenistan	Both sexes	9.5	9.7	9.9	14.4
	Females	5.6	3.4	5.4	10.5
	Males	13.2	16.3	15.7	24.5
Ukraine	Both sexes	14.5	27.0	25.8	31.1
	Females	3.7	5.5	9.1	14.2
	Males	24.7	48.6	46.5	69.6

WHO European Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
United Kingdom of Great Britain and Northern Ireland	Both sexes	10.3	14.4	11.5	8.2
	Females	4.9	6.5	5.5	4.6
	Males	15.4	22.4	18.0	12.6
Uzbekistan	Both sexes	14.3	9.2	10.3	16.0
	Females	14.4	4.2	4.9	12.5
	Males	14.2	14.1	16.0	21.8

WHO South-East Asia Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Bangladesh	Both sexes	4.7	3.2	3.1	5.2
	Females	4.0	1.7	1.7	2.5
	Males	5.5	4.8	4.4	8.2
Bhutan	Both sexes	4.5	5.1	8.4	18.8
	Females	3.4	3.1	6.2	13.1
	Males	5.4	6.8	10.4	24.6
Democratic People's Republic of Korea	Both sexes	5.2	8.4	15.5	27.1
	Females	4.0	7.3	12.8	22.6
	Males	6.3	9.4	18.3	34.4
India	Both sexes	15.9	16.2	17.6	22.2
	Females	17.0	12.6	14.3	18.0
	Males	15.0	19.5	20.8	27.1
Indonesia	Both sexes	1.4	0.8	2.0	6.4
	Females	0.6	0.7	1.7	5.4
	Males	2.2	1.0	2.4	7.8
Maldives	Both sexes	1.2	1.2	1.4	21.2
	Females	0.3	0.1	0.3	4.6
	Males	1.6	1.7	2.3	36.1

WHO South-East Asia Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Myanmar	Both sexes	3.3	3.6	4.2	7.1
	Females	1.2	1.0	1.7	2.6
	Males	5.4	6.2	6.9	13.7
Nepal	Both sexes	14.0	12.7	14.7	24.5
	Females	11.2	7.8	9.2	13.1
	Males	16.9	18.6	20.6	38.6
Sri Lanka	Both sexes	17.9	17.6	17.5	33.0
	Females	10.4	4.8	5.2	14.1
	Males	25.3	31.4	31.7	61.4
Thailand	Both sexes	21.5	26.3	12.4	13.8
	Females	5.7	7.1	5.7	7.6
	Males	36.8	45.9	20.0	22.3
Timor-Leste	Both sexes	4.9	4.8	6.1	9.1
	Females	3.5	2.4	4.0	5.3
	Males	6.3	7.2	8.1	13.6

WHO Western Pacific Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Australia	Both sexes	14.7	16.4	15.9	16.6
	Females	8.1	8.2	7.4	8.9
	Males	21.0	24.8	24.8	25.6
Brunei Darussalam	Both sexes	4.8	4.3	1.0	5.1
	Females	0.0	1.1	0.0	2.8
	Males	8.8	6.9	2.0	7.7
Cambodia	Both sexes	5.0	6.0	8.7	12.1
	Females	3.1	3.0	5.6	7.2
	Males	6.9	9.0	12.6	20.2

WHO Western Pacific Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
China	Both sexes	4.2	6.2	12.2	38.5
	Females	2.9	4.5	10.4	30.9
	Males	5.3	7.9	14.0	48.2
Fiji	Both sexes	15.0	8.5	9.6	11.9
	Females	10.8	4.1	6.3	7.9
	Males	19.3	12.8	13.1	17.7
Japan	Both sexes	18.7	20.3	20.5	18.7
	Females	13.8	12.2	13.0	12.9
	Males	23.4	28.2	27.8	26.6
Kiribati	Both sexes	31.1	27.0	16.8	16.0
	Females	9.5	4.9	5.6	7.7
	Males	53.2	51.7	30.8	30.5
Lao People's Democratic Republic	Both sexes	6.9	5.6	6.7	10.2
	Females	4.6	2.4	3.4	4.8
	Males	9.1	8.7	10.2	16.9
Malaysia	Both sexes	5.1	6.9	10.2	14.4
	Females	1.5	2.3	4.6	7.3
	Males	8.3	10.7	15.7	22.3
Micronesia (Federated States of)	Both sexes	33.4	29.5	19.3	19.9
	Females	15.2	9.3	10.4	13.7
	Males	51.2	49.8	29.8	29.4
Mongolia	Both sexes	26.2	31.4	20.4	21.0
	Females	7.5	5.9	7.0	12.4
	Males	43.9	56.5	36.6	35.2
New Zealand	Both sexes	15.2	15.4	14.1	11.7
	Females	9.4	6.7	6.6	6.0
	Males	20.6	24.4	21.9	18.3
Papua New Guinea	Both sexes	2.5	2.6	3.4	4.2
	Females	2.5	1.4	2.6	2.9
	Males	2.5	3.8	4.1	5.4

WHO Western Pacific Region					
Crude suicide rate, ^a by age group, 2021					
Country	Sex	15-29 years	30-49 years	50-69 years	70+ years
Philippines	Both sexes	4.6	4.2	5.1	13.6
	Females	2.2	1.5	2.3	11.5
	Males	6.9	6.8	8.1	17.2
Republic of Korea	Both sexes	20.2	28.0	31.3	56.5
	Females	17.2	19.0	15.7	28.8
	Males	22.8	36.4	47.1	96.8
Samoa	Both sexes	24.5	18.4	12.9	16.0
	Females	14.5	8.5	8.3	12.8
	Males	34.5	28.7	17.2	20.4
Singapore	Both sexes	9.4	5.7	11.9	13.1
	Females	7.1	2.6	6.4	8.4
	Males	11.2	8.6	17.3	19.2
Solomon Islands	Both sexes	33.9	32.8	22.3	23.5
	Females	18.3	12.2	12.9	15.5
	Males	48.9	52.6	31.1	31.8
Tonga	Both sexes	8.1	6.5	6.1	9.6
	Females	5.9	3.9	4.7	8.1
	Males	10.5	9.8	7.5	11.5
Vanuatu	Both sexes	29.7	24.7	17.0	22.1
	Females	13.7	7.0	8.8	13.2
	Males	46.2	42.2	25.1	29.1
Viet Nam	Both sexes	6.0	7.8	13.2	22.9
	Females	3.2	3.4	8.0	17.2
	Males	8.7	12.3	19.2	33.0

^a per 100 000 population

